

The Lattice of \mathcal{J} -Classes of (\mathcal{J}, σ) -Irreducible Monoids

Zhuo Li* and Lex E. Renner

*Department of Mathematics, University of Western Ontario, London,
 Ontario N6A 5B7, Canada*

Communicated by T. E. Hall

Received January 29, 1996

In this paper, we introduce a new concept, namely, the (\mathcal{J}, σ) -irreducible monoid. Let G_0 be a simple algebraic group. Let σ be a surjective endomorphism of G_0 such that $(G_0)_\sigma$, the fixed points of G_0 under σ , is finite. We construct a (\mathcal{J}, σ) -irreducible monoid M with G the unit group. Extending σ to M , we obtain a finite monoid M_σ which is \mathcal{J} -irreducible. If M is \mathcal{J} -irreducible, we give a precise recipe to determine the lattices of \mathcal{J} -classes of M and M_σ . © 1997 Academic Press

1. PRELIMINARIES

1.1. An algebraic monoid is an affine algebraic variety M defined over an algebraically closed field K together with an associative morphism $m: M \times M \rightarrow M$ and a two-sided unit $1 \in M$ for m . M is an irreducible algebraic monoid if M is irreducible as a variety. $E(M) = \{e \in M \mid e^2 = e\}$ is the set of idempotents of M . $G = G(M) = \{x \in M \mid \exists y \in M \text{ s.t. } xy = yx = 1\}$ is the unit group of M . An irreducible monoid M is reductive if its unit group G is reductive. An irreducible monoid M with 0 is semisimple if G is reductive and $\dim ZG = 1$.

(*) Throughout this paper, the monoids are always reductive.

1.2. For M and G as above, let T be some maximal torus of G and let $B \supset T$ be the Borel subgroup of G . $\Delta = \{\alpha_1, \alpha_2, \dots, \alpha_l\}$ is the fundamental root system with respect to B and T , and Φ (Φ^+ or Φ^-) is the root (positive or negative) system. $W = N_G(T)/T$ is the Weyl group, which is isomorphic to the group generated by the set of simple reflections $S = \{s_{\alpha_1}, s_{\alpha_2}, \dots, s_{\alpha_l}\}$.

* Partially supported by R. M. Kane's grant from NSERC.

1.3. Green's \mathcal{J} -relation on M is defined by $a\mathcal{J}b$ if $MaM = MbM$. The equivalence classes for this relation are called \mathcal{J} -classes. It is easy to see that $MaM = MbM$ if and only if $GaG = GbG$ and in this case the \mathcal{J} -class of a is GaG . If we consider the group $G \times G$ acting on M by $(g, h) \cdot a = gah^{-1}$ for $g, h \in G$ and $a \in M$, then the J -class of a is the orbit GaG . We define a partial order on the set of \mathcal{J} -classes by $MaM \leq MbM$ iff $MaM \subseteq MbM$. Define $\mathcal{U}(M) = \{J \text{ some } \mathcal{J}\text{-class} \mid E(J) \neq \emptyset\}$. $\mathcal{U}(M)$ is the set of \mathcal{J} -classes of M which contain idempotents. It inherits a partial order from the set of all \mathcal{J} -classes. By the assumption of $(*)$, it follows that $\mathcal{U}(M)$ equals the set of all \mathcal{J} -classes on M [11, Proposition 3.6]. From the semigroup theoretic point of view, one interesting problem is to describe $\mathcal{U}(M)$. By [5, Corollaries 6.8 and 6.10, Theorem 6.25] the problem of determining $\mathcal{U}(M)$ is to find the set of W -conjugacy classes in $E(\bar{T})$. Note that $E(\bar{T})$ is a finite, relatively complemented lattice with partial order $e < f$ defined by $ef = fe = e \neq f$ [5, Theorem 6.20].

1.4. Let M be an irreducible monoid with unit group G . Then $\Lambda \subseteq E(\bar{T})$ is a *cross-section lattice* if

- (1) $|\Lambda \cap J| = 1$ for all $J \in \mathcal{U}(M)$.
- (2) If $e, f \in \Lambda$, then $J_e \geq J_f$ implies $e \geq f$.

The cross-section lattice is a sublattice of $E(\bar{T})$, and $\Lambda = \{e \in E(\bar{T}) \mid B \subseteq C_G^r(e)\}$ [5, Theorem 9.10]. $C_G^r = \{x \in G \mid ge = ege\}$ is the *right centralizer* of e . So finding $\mathcal{U}(M)$ is equivalent to finding Λ .

2. \mathcal{J} -IRREDUCIBLE MONOIDS

2.1. A reductive monoid M is a \mathcal{J} -irreducible monoid if all minimal nonzero idempotents are conjugate.

To construct a \mathcal{J} -irreducible monoid, we choose an algebraic group G_0 and a rational representation $\rho: G_0 \rightarrow GL_n$ for some n , with finite kernel. Then $M(\rho) = \overline{K^*\rho(G_0)}$ is an irreducible monoid but not necessarily \mathcal{J} -irreducible. However, $M(\rho)$ is \mathcal{J} -irreducible if ρ is an irreducible representation and G_0 is semisimple (see [8, Corollary 8.3.3]). Putcha and Renner [6, 4.16] gave a precise recipe for constructing the lattice of \mathcal{J} -classes and the type map of a \mathcal{J} -irreducible monoid M . We sketch the main results here. For $e \in \Lambda$, let

$$I(e) = \{\alpha \in \Delta \mid s_\alpha e = es_\alpha \neq e\}$$

and

$$J(e) = \{\alpha \in \Delta \mid s_\alpha e = es_\alpha = e\}.$$

Let e_0 be the unique minimal nonzero idempotent in Λ , and define $J_0 = J(e_0)$.

THEOREM 2.2 (Putcha and Renner [6, 4.11, 4.12, 4.16]). (1) $J(e) = \{\alpha \in J_0 \setminus I(e) \mid s_\alpha s_\beta = s_\beta s_\alpha \text{ for all } \beta \in I(e)\}$.

(2) $I: \Lambda \rightarrow \mathcal{P}(\Delta)$, $e \rightarrow I(e)$, is an order-preserving injection.

(3) $S(\subseteq \Delta) = I(e)$ for some $e \in \Lambda$ iff no component of S lies entirely in J_0 .

PROPOSITION 2.3. Let ρ be the irreducible representation corresponding to a dominant weight $\mu = \sum_{i=1}^l a_i \mu_i$. Let $I(\mu) = \{\alpha_i \mid a_i \neq 0\}$. Then $J_0 = \Delta \setminus I(\mu)$.

Proof. Let $v \in V$ be the highest weight vector of weight μ . Then, by [2, Section 31], the isotropic group of Kv is $P_L = BW_L B$, where $L = \{\alpha \in \Delta \mid \rho(s_\alpha)v = v\} = \Delta \setminus I(\mu)$. So it suffices to show that $L = J_0$. Let Γ be the set of weights of the representation ρ . Let $V_\lambda = \{v \in V \mid \rho(t)v = \langle \lambda, t \rangle v = \lambda(t)v\}$ for $\lambda \in \Gamma$. Then $V = \bigoplus_{\lambda \in \Gamma} V_\lambda$. Note that $V_\mu = Kv$. To simplify the notation, we denote $\rho(g)x$ by $g \cdot x$ for $g \in G$ and $x \in V$. Now, by [6, Theorem 5.3], $Kv = e_0 \cdot V$, since $Kv = V^{B_u}$. So

$$e_0 \cdot v = \langle \mu, e_0 \rangle v = v,$$

since $\langle \mu, e_0 \rangle \neq 0$ and e_0 is an idempotent. Also

$$e_0 \cdot v_\lambda = 0 \quad \text{for } v_\lambda \in V_\lambda \text{ and } \mu \neq \lambda \in \Gamma.$$

Then

$$\begin{aligned} e_0 \cdot v &= e_0 s_\alpha \cdot v = s_\alpha s_\alpha^{-1} e_0 s_\alpha \cdot v \\ &= s_\alpha \cdot \langle \mu, s_\alpha^{-1} e_0 s_\alpha \rangle v \\ &= s_\alpha \cdot \langle s_\alpha \mu, e_0 \rangle v = s_\alpha \cdot \langle \mu, e_0 \rangle v \\ &= s_\alpha e_0 \cdot v \quad \text{for } \alpha \in L, \end{aligned}$$

and

$$\begin{aligned} 0 &= e_0 \cdot v_\lambda = s_\alpha e_0 \cdot v_\lambda \\ e_0 s_\alpha \cdot v_\lambda &= s_\alpha s_\alpha^{-1} e_0 s_\alpha \cdot v_\lambda \\ &= s_\alpha \cdot \langle \lambda, (s_\alpha^{-1} e_0 s_\alpha) \rangle v_\lambda \\ &= s_\alpha \cdot \langle s_\alpha \lambda, e_0 \rangle v_\lambda \\ &= 0 \quad \text{for } \alpha \in L \text{ and } \mu \neq \lambda \in \Gamma, \end{aligned}$$

since $s_\alpha \lambda < \mu$. Thus $e_0 = s_\alpha e_0 = e_0 s_\alpha$, so that $L \subseteq J_0$.

Conversely, for $\alpha \in J_0$,

$$v = e_0 \cdot v = s_\alpha e_0 \cdot v = e_0 s_\alpha \cdot v.$$

Then

$$s_\alpha \cdot v = v.$$

Thus $\alpha \in L$, so that $J_0 \subseteq L$. ■

3. IDEMPOTENTS AND CONVEX HULL

3.1. Using the theory of torus embedding, Solomon [11, Section 5] gives a detailed description about how to compute idempotents in $E(\bar{T})$ and the cross-section lattices Λ . The following result is due to the second named author (see [11, Section 5] and [8, Section 3]).

PROPOSITION 3.2. *Let M be the monoid as constructed in 2.1 by a rational representation ρ . Let $C^\vee(1)$ be a convex hull of the set $\Phi(\rho)$ of weights of ρ . Then*

- (1) *There is a lattice isomorphism from the face lattice of $\mathcal{A}(C^\vee(1))$, including the empty face, to $E(\bar{T})$.*
- (2) *There is a lattice isomorphism from the set of W -orbits of $\mathcal{A}(C^\vee(1))$ to the cross-section lattice $\Lambda \setminus 0$.*
- (3) *If ρ corresponds to a dominant weight μ , then $C^\vee(1)$ is the convex hull of $W \cdot \mu$.*

4. (\mathcal{J}, σ) -IRREDUCIBLE MONOIDS

4.1. A (\mathcal{J}, σ) -irreducible monoid M is a reductive monoid with an endomorphism σ of M such that (W, σ) acts transitively on $E_1(\bar{T})$, the set of minimal nonzero idempotents in $E(\bar{T})$. More precisely, let $\Lambda_1 = \Lambda \cap E_1(\bar{T})$. Then σ permutes the set Λ_1 . In other words, for any $e \in E_1(\bar{T})$,

$$E_1(\bar{T}) = W \cdot e \cup W \cdot \sigma(e) \cup W \cdot \sigma^2(e) \dots$$

Obviously, M is a (\mathcal{J}, σ) -irreducible monoid if M is \mathcal{J} -irreducible and $\sigma = 1$. This special case is well studied in [6]. However, most of the (\mathcal{J}, σ) -irreducible monoids are usually not \mathcal{J} -irreducible.

The definition of a (\mathcal{J}, σ) -irreducible monoid is motivated by the second named author's construction of finite reductive monoids as in [7, Section 4] or [9].

Next we construct a class of (\mathcal{J}, σ) -irreducible monoids such that $M_\sigma = \{x \in M \mid \sigma(x) = x\}$ is finite. Let G_0 be a simple algebraic group. Chevalley [12, Section 11] classified all endomorphisms $\sigma: G_0 \rightarrow G_0$ such

that $G_{0\sigma} = \{x \in G_0 \mid \sigma(x) = x\}$ is a finite group. Let $\sigma: G_0 \rightarrow G_0$ be a surjective morphism such that $G_{0\sigma}$ is finite. Then there is a Borel subgroup B_0 of G_0 and a maximal torus T_0 of B_0 such that $\sigma(T_0) = T_0$ and $\sigma(B_0) = B_0$ [12, Section 10]. Let Δ be the fundamental root system associated with the maximal torus T_0 of G_0 . Let $\sigma^*: X(T_0) \rightarrow X(T_0)$ be the map induced by $\sigma|_{T_0}$. Then there exists a nontrivial symmetric automorphism of the Dynkin diagram ϱ and a scalar factor $q: \Phi \rightarrow \mathbf{N}$ such that

$$\sigma^*(\varrho(\alpha)) = q(\alpha)\alpha \quad \text{for all } \alpha \in \Delta. \quad (4.1.1)$$

There are two types of σ .

(1) *Chevalley and Steinberg type*: If the root systems are A_n , D_n , and E_6 , then $\sigma = Fr_q \circ \varrho$, where $Fr_q(t) = t^q$ for all $t \in T$ and q is some power of p , the characteristic of the field K .

(2) *Suzuki and Ree type*: If the root system is C_2 , G_2 , or F_4 , then $q(\alpha) = p^a$ for α long, and $q(\alpha) = p^{a+1}$ for α short. Here $p = 2$ for C_2 and F_4 , and $p = 3$ for G_2 .

4.2. Suppose σ is of Steinberg type. Choose a dominant weight μ in $X(T)$. Let $\rho_i: G_0 \rightarrow GL(V_i)$ be the irreducible representations corresponding to $\varrho^{i-1}(\mu)$ for $i = 1, 2$ (or 3 for D_4 only).

Define

$$\rho = \bigoplus \rho_i: G_0 \rightarrow GL\left(\bigoplus V_i\right) \quad (4.2.1)$$

and $M = M(\rho) = \overline{K^*\rho(G_0)}$.

THEOREM 4.3. (1) $M(\rho)$ is a (\mathcal{J}, σ) -irreducible monoid.

(2) M_σ is finite.

(3) M_σ is \mathcal{J} -irreducible.

We leave the proof for later. Let $T = K^*\rho(T_0)$ and $G = K^*\rho(G_0)$. Since ρ has a finite kernel, $\rho(G_0)$ is also a simple algebraic group. We can identify the root system of $\rho(G_0)$ relative to $\rho(T_0)$ and $\rho(B_0)$ with that of G_0 . Also there is a commutative diagram:

$$\begin{array}{ccc} G_0 & \xrightarrow{\sigma} & G_0 \\ \rho \downarrow & & \downarrow \rho \\ \rho(G_0) & \longrightarrow & \rho(G_0) \end{array} \quad (4.3.1)$$

We use the same notation σ for the bottom map.

Since $X(T) = X(\rho(T_0)) \oplus \mathbf{Z}$, let $(\mu, 1) \in X(T)$ and σ^* extends to $X(T)$ by $\sigma_1^*(\chi, t) = (\sigma^*(\chi), t^q)$. Then σ_1^* is induced from $\sigma_1: G \rightarrow G$, $\sigma_1(g, t) = (\sigma(g), t^q)$. Note that $\sigma^*(\mu) = q\mu_2$.

Let C_i^\vee be the smallest convex cone containing $(W \cdot \varrho^{i-1}(\mu), 1)$. Let C^\vee be the smallest convex cone containing $\bigcup C_i^\vee$. Then $M(\rho)$ is the semisimple monoid corresponding to the *polyhedral root system* $(\Phi, X(T), C^\vee = X(\bar{T}))$ (see [8]).

LEMMA 4.4 (Renner [10]). $\sigma_1: G \rightarrow G$ extends to a surjective morphism $\sigma_1: M \rightarrow M$ such that G_{σ_1} and M_{σ_1} are finite iff

$$(1) \quad \sigma_1^*(C^\vee) \subseteq C^\vee, \text{ and}$$

$$(3) \quad nC^\vee \subseteq \sigma_1^*(C^\vee).$$

4.5. *Proof of Theorem 4.3.* For (1), it is clearly true by the construction.

For (2), it follows since C^\vee satisfies (1) and (2) of Lemma 4.4. Changing notation, let $\sigma = \sigma_1$ denote the finite morphism $\sigma: G \rightarrow G$ and $\sigma: M \rightarrow M$ such that G_σ and M_σ are finite. So M_σ is the finite monoid with unit group G_σ .

For (3), it is enough to show the case $\varrho^2 = 1$. Let e_1, e_2 be two nonzero minimal idempotents in the cross-section lattice Λ of M . Then $\sigma(e_1) = e_2$ and $\sigma(e_2) = e_1$. So

$$\sigma(e_1 \vee e_2) = \sigma(e_1) \vee \sigma(e_2) = e_1 \vee e_2 \in \Lambda_\sigma, \quad (4.5.1)$$

where $\Lambda_\sigma = \{e \in \Lambda \mid \sigma(e) = e\}$ is the cross-section lattice of M_σ . For any $e \in \Lambda_\sigma \setminus 0$, either $e > e_1$ or $e > e_2$. In the former case, $e = \sigma(e) > \sigma(e_1) = e_2$, so that $e > e_1 \vee e_2$. Then $e_1 \vee e_2$ is the unique nonzero minimal idempotent in Λ_σ . ■

4.6. For σ of Suzuki or Ree type, one cannot obtain a finite monoid M_σ from a semisimple monoid M as in [7, Section 4], because the requirement that $\sigma: M \rightarrow M$ would imply that $\sigma^2(t) = t^{p^{2a+1}}$ for all $t \in Z(G(M))^0$ and some $a > 0$ [12]. But that $\sigma(t) = t^m$ implies $\sigma^2(t) = t^{m^2} = t^{p^{2a+1}}$ for all t . That is impossible. The following construction is contained in [7].

Assume G_0 is a simple algebraic group of type C_2 , F_4 , or G_2 , and $\sigma: G_0 \rightarrow G_0$ is an endomorphism of Ree or Suzuki type. Assuming $\sigma(T_0) = T_0$ and $\sigma(B_0) = B_0$ with $T_0 \subseteq B_0$, we can arrange the simple roots $\Delta = \{\alpha_i, \beta_i\}_{i=1}^s$ ($s = 1$ or 2) for α_i short and β_i long so that $\sigma^*(\alpha_i) = p^q \beta_i$ and $\sigma^*(\beta_i) = p^{a+1} \alpha_i$. Choose $\mu = \sum_{i=1}^{2s} a_i \mu_i$ (some a_i may be 0, not as in [7]), a dominant weight. Then $\sigma^*(\mu)$ is also a dominant weight, and $\sigma^*(\mu)$ is not a multiple of μ since σ^* has no rational eigenvalues on $X(T) \otimes \mathbf{R}$. Let $\mu' = \sigma^*(\mu)/p^a$. Let $\rho_1: G \rightarrow GL(V_1)$ and $\rho_2: G \rightarrow GL(V_2)$ be the irreducible representations with highest weight μ and μ' , respectively, and let $\rho = \rho_1 \oplus \rho_2$. Define

$$M \text{ to be the normalization of } \overline{\rho(G_0)(K^* \times K^*)}.$$

Let $G = \overline{\rho(G_0)(K^* \times K^*)}$, and let T be the maximal torus of G containing T_0 . Then $X(T) = X(T_0) \oplus \mathbb{Z}^2$. Let $\chi = (\mu, 1, 0)$ and $\chi' = (\mu', 0, 1)$. Then $C^\vee = X(\bar{T})$ is the smallest polyhedral cone containing $W \cdot \chi \cup W \cdot \chi'$. The morphism $\sigma: G_0 \rightarrow G_0$ extends to $\sigma: G \rightarrow G$ by $\sigma(g, s, t) = (\sigma(g), t^{p^{a+1}}, s^{p^a})$ and this yields $\sigma^*: X(T) \rightarrow X(T)$ by $\sigma^*(\gamma, \alpha, \beta) = (\sigma^*(\gamma), p^{a+1}\beta, p^a\alpha)$. Since $\sigma^*(\chi) = p^a\chi'$ and $\sigma^*(\chi') = p^{a+1}\chi$, $\sigma^*(X(\bar{T})) \subseteq X(\bar{T})$ with $p^{2a+1}X(\bar{T}) \subseteq \sigma^*X(\bar{T})$. Thus there exists a unique $\sigma: M \rightarrow M$ extending $\sigma: \bar{T} \rightarrow \bar{T}$ and $\sigma: G \rightarrow G$ by Lemma 4.4.

PROPOSITION 4.7 [7, Theorem 4.10]. $M(\rho)$ is (\mathcal{J}, σ) -irreducible and M_σ is finite and \mathcal{J} -irreducible.

4.8. We are now in a position to ask the following questions.

- (A) What is $\mathcal{U}(M)$ or, equivalently, Λ ?
- (B) What is $\mathcal{U}(M_\sigma)$ or, equivalently, Λ_σ ?

If $\mu = \sum_{i=1}^l a_i \mu_i$, define $I(\mu) = \{\alpha_i \in \Delta \mid a_i \neq 0\}$. More precisely, we should answer these two questions in terms of the following separate cases:

- (1) $\sigma^*(\mu) = q^n(\mu)$.
- (2) $\sigma^*(\mu) \neq q^n(\mu)$.
 - (a) $I(\mu) = I(\sigma^*(\mu))$.
 - (b) $I(\mu) \neq I(\sigma^*(\mu))$.

PROPOSITION 4.9 ([11, 5.49], [6, Section 4], and [8, 3.5]). For Steinberg type, let $C_i^\vee(1)$ be the convex hull of $W \cdot \rho^{i-1}\mu$. Let $C^\vee(1)$ be the convex hull of $\bigcup C_i^\vee(1)$. For Suzuki or Ree type, let $C_1^\vee(1)$ be the convex hull of $W \cdot (\mu, 1, 0)$. Let $C_2^\vee(1)$ be the convex hull of $W \cdot (\mu', 0, 1)$. Let $C^\vee(1)$ be the convex hull of $C_1^\vee(1)$ and $C_2^\vee(1)$. Then

- (1) There is a lattice isomorphism from the face lattice of $\mathcal{A}(C^\vee(1))$, including the empty face, to $E(\bar{T})$.
- (2) There is a lattice isomorphism from the set of W -orbits of $\mathcal{A}(C^\vee(1))$ to the cross-section lattice $\Lambda \setminus 0$.

COROLLARY 4.10. If σ is of Steinberg type and $\sigma^*(\mu) = q(\mu)$, then M is \mathcal{J} -irreducible and $\mathcal{U}(M) \cong \mathcal{U}(M(\rho_1))$, where $M(\rho_1) = \overline{K^*\rho_1(G_0)}$.

Proof. This follows immediately from (2) of proposition 4.9 and the fact that $C^\vee(1) = C_i^\vee(1)$ for all i . ■

Let Δ_ϱ be the set of equivalence classes under ϱ , i.e., the set of ϱ -orbits of Δ . It can be considered as the fundamental root system relative to B_σ and T_σ [1, Section 13]. More precisely, we identify Δ_ϱ case by case as

follows:

- (1) For case A_l , let $l = 2k - 1$ and $\Delta = \{\alpha_1, \dots, \alpha_l\}$. Then Δ_ϱ is

$$(\alpha_1 + \alpha_{2k-1})/2, \dots, (\alpha_{k-1} + \alpha_{k+1})/2, \alpha_k$$

of type C_k .

- (2) For case A_l , let $l = 2k$ and $\Delta = \{\alpha_1, \dots, \alpha_l\}$. Then Δ_ϱ is

$$(\alpha_1 + \alpha_{2k})/2, \dots, (\alpha_k + \alpha_{k+1})/2$$

of type B_k .

- (3) For case D_l , let $\Delta = \{\alpha_1, \dots, \alpha_l\}$ and $\varrho(\alpha_{l-1}) = \alpha_l$. Then Δ_ϱ is

$$\alpha_1, \alpha_2, \dots, \alpha_{l-2}, (\alpha_{l-1} + \alpha_l)/2$$

of type B_{l-1} .

- (4) For case D_4 , let $\Delta = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$. Let $\varrho^3 = 1$ and let α_1 be the root in the center of the Dynkin diagram of D_4 . Then Δ_ϱ is

$$\alpha_1, (\alpha_2 + \alpha_3 + \alpha_4)/3$$

of type G_2 .

- (5) For case E_6 , let $\Delta = \{\alpha_1, \dots, \alpha_6\}$. Let $\varrho(\alpha_i) = \alpha_{7-i}$ for $i = 1, 2$. Then Δ_ϱ is

$$(\alpha_1 + \alpha_6)/2, (\alpha_2 + \alpha_5)/2, \alpha_3, \alpha_4$$

of type F_4 .

Define a natural projection $\pi: \Delta \rightarrow \Delta_\varrho$ by

$$\pi(\alpha) = \frac{\sum_{i=1}^m \rho^{i-1}(\mu)}{m},$$

where m is the order of ϱ . Let μ and σ be as in Corollary 4.10. Let $I'(\mu) = \pi(I(\mu))$. Since $\varrho(I(\mu)) = I(\mu)$, then $\pi^{-1}(I'(\mu)) = I(\mu)$. Let $J'_0 = \Delta_\varrho \setminus I'(\mu)$. Since $\varrho(I(\mu)) = I(\mu)$, then $\pi^{-1}(I'(\mu)) = I(\mu)$. Thus $J'_0 = \pi(J_0)$. We have

THEOREM 4.11. (1) *There is an order-preserving injection*

$$e \in \Lambda_\sigma \rightarrow K(e) \subseteq \mathcal{P}(\Delta_\varrho).$$

(2) *Any $S(\subseteq \Delta_\varrho) = K(e)$ for some $e \in \Lambda_\sigma$ if and only if S has no connected component completely contained in J'_0 .*

Proof. (1) For $e \in \Lambda$, according to Theorem 2.2, there is a subset $I(e) \subseteq \Delta$ such that no connected components of $I(e)$ lie in J_0 by Theorem 2.2. Furthermore, $e \in \Lambda_\sigma$ if and only if $\varrho(I(e)) = I(e)$. Let $K(e) = \pi(I(e))$. This proves (1).

(2) The above proof implies the “if” part. So it suffices to show the “only if” part. That is, if $S(\subseteq \Delta_\varrho)$ has no connected components contained in J'_0 , then $S = K(e)$ for some $e \in \Lambda_\sigma$.


Let $I = \pi^{-1}(S)$. Then $\varrho I = I$ by the definition of π . It is enough to show that I has no connected components contained in J_0 .

Suppose that I has a connected component I_1 contained in J_0 . Then $\varrho(I_1)$ is also a connected component of I , and $\varrho(I_1) \subseteq J_0$ since $\varrho(J_0) = J_0$. Thus $S_1 = \pi(I_1) \subseteq \pi(J_0) = J'_0$, so that S_1 being a connected component of S is contained in J'_0 . But that is a contradiction. ■


5. CASE E_6^2

5.1. Using Theorems 2.2 and 4.11, we give a complete list of the Hasse diagrams of $\Lambda \setminus 0$ and $\Lambda_\sigma \setminus 0$ for the \mathcal{J} -irreducible monoids M and M_σ , respectively, associated with type E_6^2 .

5.2. Let the Dynkin diagram of E_6 be as follows:



Then the Dynkin diagram for Δ_ϱ is



where $\beta_4 = \alpha_4$, $\beta_3 = \alpha_3$, $\beta_2 = (\alpha_2 + \alpha_5)/2$, $\beta_1 = (\alpha_1 + \alpha_6)/2$. We assume that k, k_i for $i = 1, 2, \dots$ are positive integers in the following.

(1) $\mu = k\mu_4$. In this case $J_0 = \{\alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_6\}$ and $J_0^* = \{\beta_1, \beta_2, \beta_3\}$. $\Lambda \setminus 0$ is shown in Fig. 1, where $i_1 i_2 \dots$ represents the subset $\{\alpha_{i_1}, \alpha_{i_2}, \dots\}$ of Δ in the Hasse diagrams of $\Lambda \setminus 0$, and $\overline{i_1 i_2 \dots}$ represents the subset $\{\alpha_{i_1}, \alpha_{i_2}, \dots\}$ of Δ fixed by ϱ which corresponds to the element in the Hasse diagram of $\Lambda_\sigma \setminus 0$, henceforth. So $\Lambda_\sigma \setminus 0$ is shown in Fig. 2, where $i_1 i_2 \dots$, which equals $\pi(\overline{j_1 j_2 \dots})$ for some $\overline{j_1 j_2 \dots}$ in the Hasse diagram of $\Lambda \setminus 0$, represents the subset $\{\beta_{i_1}, \beta_{i_2}, \dots\}$ of Δ_ϱ in the Hasse diagram of $\Lambda_\sigma \setminus 0$, henceforth.

(2) $\mu = k\mu_3$. In this case $J_0 = \{\alpha_1, \alpha_2, \alpha_4, \alpha_5, \alpha_6\}$ and $J'_0 = \{\beta_1, \beta_2, \beta_4\}$. $\Lambda \setminus 0$ is shown in Fig. 3 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 4.




FIGURE 1




FIGURE 2




FIGURE 3

(3) $\mu = k_1 \mu_3 + k_2 \mu_4$. In this case $J_0 = \{\alpha_1, \alpha_2, \alpha_5, \alpha_6\}$ and $J'_0 = \{\beta_1, \beta_2\}$. $\Lambda \setminus 0$ is shown in Fig. 5 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 6.

(4) $\mu = k \mu_5 + k \mu_2$. In this case $J_0 = \{\alpha_1, \alpha_3, \alpha_4, \alpha_6\}$ and $J'_0 = \{\beta_1, \beta_3, \beta_4\}$. $\Lambda \setminus 0$ is shown in Fig. 7 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 8.




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7

(5) $\mu = k\mu_1 + k\mu_6$. In this case $J_0 = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5\}$ and $J'_0 = \{\beta_2, \beta_3, \beta_4\}$. $\Lambda \setminus 0$ is shown in Fig. 9 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 10.

(6) $\mu = k_1\mu_3 + k\mu_2 + k\mu_5$. In this case $J_0 = \{\alpha_1, \alpha_4, \alpha_6\}$ and $J'_0 = \{\beta_1, \beta_4\}$. $\Lambda \setminus 0$ is shown in Fig. 11 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 12.

(7) $\mu = k_1\mu_4 + k\mu_2 + k\mu_5$. In this case $J_0 = \{\alpha_1, \alpha_3, \alpha_6\}$ and $J'_0 = \{\beta_1, \beta_3\}$. $\Lambda \setminus 0$ is shown in Fig. 13 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 14.

(8) $\mu = k_1\mu_3 + k\mu_1 + k\mu_6$. In this case $J_0 = \{\alpha_2, \alpha_4, \alpha_5\}$ and $J'_0 = \{\beta_2, \beta_4\}$. $\Lambda \setminus 0$ is shown in Fig. 15 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 16.




FIGURE 8




FIGURE 9

(9) $\mu = k_1 \mu_4 + k \mu_1 + k \mu_6$. In this case $J_0 = \{\alpha_2, \alpha_3, \alpha_5\}$ and $J'_0 = \{\beta_2, \beta_3\}$. $\Lambda \setminus 0$ is shown in Fig. 17 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 18.

(10) $\mu = k_1 \mu_3 + k_2 \mu_4 + k \mu_1 + k \mu_6$. In this case $J_0 = \{\alpha_2, \alpha_5\}$ and $J'_0 = \{\beta_2\}$. $\Lambda \setminus 0$ is shown in Fig. 19 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 20.

(11) $\mu = k_1 \mu_3 + k_2 \mu_4 + k \mu_2 + k \mu_5$. In this case $J_0 = \{\alpha_1, \alpha_6\}$ and $J'_0 = \{\beta_1\}$. $\Lambda \setminus 0$ is shown in Fig. 21 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 22.




FIGURE 10




FIGURE 11

(12) $\mu = k_1 \mu_1 + k_1 \mu_6 + k \mu_2 + k \mu_5$. In this case $J_0 = \{\alpha_3, \alpha_4\}$ and $J'_0 = \{\beta_3, \beta_4\}$. $\Lambda \setminus 0$ is shown in Fig. 23 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 24.

(13) $\mu = k_1 \mu_1 + k_1 \mu_6 + k_2 \mu_2 + k_2 \mu_5 + k \mu_3$. In this case $J_0 = \{\alpha_4\}$ and $J'_0 = \{\beta_4\}$. $\Lambda \setminus 0$ is shown in Fig. 25 and $\Lambda_\sigma \setminus 0$ is shown in Fig. 26.




FIGURE 12




FIGURE 13

(14) $\mu = k_1 \mu_1 + k_1 \mu_6 + k_2 \mu_2 + k_2 \mu_5 + k \mu_4$. In this case $J_0\{\alpha_3\}$ and $J'_0 = \{\beta_3\}$. $\Lambda \setminus 0$ is shown in Fig. 27 and $\Delta_\sigma \setminus 0$ is shown in Fig. 28.

(15) $\mu = k_1 \mu_1 + k_1 \mu_6 + k_2 \mu_2 + k_2 \mu_5 + k_3 \mu_3 + k_4 \mu_4$. In this case $J_0 = \emptyset$ and $J'_0 = \emptyset$. So $\Lambda \setminus 0 \cong \mathcal{P}(\Delta)$ and $\Delta_\sigma \setminus 0 \cong \mathcal{P}(\Delta_\sigma)$.




FIGURE 14




FIGURE 15




FIGURE 16




FIGURE 17




FIGURE 18




FIGURE 19




FIGURE 20




FIGURE 21




FIGURE 22




FIGURE 23

Remark. Theorem 2.2 contains a precise recipe to determine the structure of the cross section lattice for a \mathcal{J} -irreducible monoid. That recipe and Theorem 4.11 give a complete solution to (1) of 4.8. It is interesting but perhaps difficult to find a general answer for (2) of 4.8. Since M_σ is \mathcal{J} -irreducible we may expect a general recipe to determine the cross section lattice at least for Λ_σ .




FIGURE 24




FIGURE 25




FIGURE 26




FIGURE 27




FIGURE 28

REFERENCES

1. R. W. Carter, "Simple Groups of Lie Type," Wiley, London/New York, 1972.
2. J. E. Humphreys, "Linear Algebraic Groups," Graduate Texts in Mathematics, Vol. 21, Springer-Verlag, New York, 1981.
3. G. Kempf, F. Knudsen, F. Mumford, and B. Saint-Donat, "Toroidal Embeddings, I" Vol. 1, Springer-Verlag, New York, 1973.
4. T. Oda, "Torus Embeddings and Applications," Tata Press, Bombay, 1978.
5. M. S. Putcha, "Linear Algebraic Monoids," London Math. Soc. Lecture Note Series 133, Cambridge Univ. Press, Cambridge, 1988.
6. M. S. Putcha and L. E. Renner, The system of idempotents and the lattice of \mathcal{J} -classes of reductive algebraic monoids, *J. Algebra* **116** (1988), 385–399.
7. M. S. Putcha and L. E. Renner, The canonical compactification of a finite group of Lie type, *Trans. Amer. Math. Soc.* **337** (1993), 305–319.
8. L. E. Renner, Classification of semisimple algebraic monoids, *Trans. Amer. Math. Soc.* **292** (1985), 193–223.
9. L. E. Renner, Finite monoids of Lie type, in "Monoids and Semigroups with Applications," pp. 278–287, World Scientific, Singapore, 1991.
10. L. E. Renner, Finite reductive monoids, in "Semigroups, Formal Languages and Groups," (J. Fountain, Ed.), pp. 369–380, Kluwer, Dordrecht, 1995.
11. L. Solomon, An introduction to reductive monoids, in "Proceedings of Groups, Semigroups and Formal Languages," 1993, pp. 295–352.
12. R. Steinberg, Endomorphisms of linear algebraic groups, *Mem. Amer. Math. Soc.* **80** (1968).